Ciflocin

Ciprofloxacin USP

Description

Ciprofloxacin is a synthetic fluoro-quinolone derivative. It has anti-bacterial activity against a wide range of gram-positive & gram negative bacteria.

Composition

Ciflocin 500 mg Tablet: Each film coated tablet contains Ciprofloxacin hydrochloride USP 500 mg.

Indications

Ciprofloxacin indicated for UTI, RTI, Infectious diarrhoea, infections of ear, nose, throat, eyes, SSTI, infections of bone & joints, gonorrhea, severe systemic infections & gram negative pneumonia.

Dosage & Administration

Ciflocin Tablet: The dose range for adults is 250 mg to 750 mg twice daily or as advised physician and for children: 10-30 mg/kg/day, or as advised physician.

Side Effects

Possible side effects arise with Ciprofloxacin administration like nausea, vomiting, diarrhoea, dyspepsia, skin rash, headache, dizziness, confusions.

Contraindications

Ciprofloxacin contraindicated in patients who are hypersensitive to Ciprofloxacin or any member of the quinolone group of drugs, growing adolescents, pregnancy & lactation, epilepsy and history of CNS disorder.

Drug Interactions

Ciprofloxacin should be given with care to patients taking other drug that effect the Liver & Kidney.

Precautions and Warnings

It should be used with caution in patients with suspected or known CNS disorders such as arteriosclerosis or epilepsy or other factors which predispose to seizures and convulsion. Ciprofloxacin may be taken with or without meals and to drink fluids liberally. Concurrent administration of Ciprofloxacin should be avoided with magnesium/aluminium antacids, or sucralfate or with other products containing calcium, iron and zinc. These products may be taken two hours after or six hours before Ciprofloxacin. Ciprofloxacin should not be taken concurrently with milk or yogurt alone, since absorption of Ciprofloxacin may be significantly reduced. Dietary calcium is a part of a meal, however, does not significantly affect the Ciprofloxacin absorption.

Storage

Store below 30^{o}C, protected from light & moisture. Keep out of reach of children.

How Supplied

Ciflocin 500 mg Tablet: Box containing 3 x 10 tablets in blister pack.

Manufactured by:

Astra Biopharmaceuticals Ltd.
Gouripur, Ashulia, Dhaka, Bangladesh