Deslorat

Desloratadine

Description:
Desloratat is a preparation of Desloratadine. Desloratadine is a long-acting tricyclic histamine antagonist with selective H1-receptor antagonist activity. It is a non-sedating antihistamine. Desloratadine has effects on the chronic inflammatory response to allergens.

Composition:
Deslorat Tablet: Each film-coated tablet contains Desloratadine USP 5 mg.

Indication:
Deslorat is indicated for the relief of symptoms associated with seasonal and perennial allergic rhinitis, such as sneezing, nasal discharge & itching, congestion/stuffiness, as well as ocular itching, tearing and redness, itching of palate and coughing.
Deslorat is also indicated for the relief of symptoms associated with chronic idiopathic urticaria such as the relief of itching and the size & number of hives.

Dosage & Administration:
Deslorat Tablet:
Adults and children 12 years of age and over: 1 tablet daily
In adult patients with liver or renal impairment: A starting dose of one 5 mg tablet every other day is recommended based on pharmacokinetic data.

Side Effects:
Desloratadine is generally well tolerated. However, dry mouth, fatigue, somnolence and myalgia are commonly reported side-effects. Less common side-effects may include dizziness, headache and nausea. Rarely rash, pruritus and urticaria may occur.

Contraindications:
Desloratadine tablets are contraindicated in patients who are hypersensitive to this medication or to any of its ingredients or to loratadine

Over Dose:
No clinically relevant adverse events have been reported in case of overdosage. However, in the event of overdosage, symptomatic and supportive treatment is recommended.

Precautions:
Pregnancy Category C: Desloratadine was not teratogenic in rats or rabbit. There are no adequate and well controlled studies of Desloratadine in pregnant women. Because animal reproduction studies are not always predictive of human response. Desloratadine should be used during pregnancy only if clearly needed. Nursing Mothers: Desloratadine passes into breast milk, therefore a decision should be made whether to discontinue nursing or to discontinue Desloratadine, taking into account the importance of the drug to the mother. Geriatric Use: In general, dose selection for an elderly patient should be cautious, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and of concomitant disease or other drug therapy.

Storage:
Store below 30^{o}C, protect from light & moisture. Keep all medicine out from children's reach.

How Supplied:
Deslorat Tablet: Each box contains 100 tablets in blister pack.

Manufactured by:
Astra Biopharmaceuticals Ltd.
Gouripur, Ashulia, Dhaka, Bangladesh