

Moov-SR

Diclofenac Sodium USP

Moov SR Capsule

Each capsule contains Diclofenac Sodium USP 100 mg in sustained release formulation.

Pharmacology

Diclofenac is a potent NSAID having both strong anti-inflammatory & analgesic properties. It inhibits synthesis of prostaglandin from arachidonic acid by blocking the cyclooxygenase enzyme.

Indications

Rheumatology: Inflammatory and degenerative forms of rheumatism, chronic involutive, polyarthritis, ankylosing spondylarthritis, osteoarthritis, spondylarthroses, acute gout, peri-articular rheumatic disorders. Surgery and Traumatology: Sprain, bruises, dislocations, fractures, soft-tissue injuries, surgical interventions. Obstetrics and Gynecology: Primary dysmenorrhoea, episiotomy, adnexitis, endometritis, parametritis, salpingitis, and mastitis. Otorhinolaryngology: As pre-operative medication for the prevention of pain, inflammation, and swelling. Dentistry: Post-operative and post-traumatic pain, inflammation, and swelling. Other indications: For the prevention of pain and treatment of inflammation and swelling of patients operated in the urogenital tract, renal and biliary colic.

Dosage & Administrations

Moov SR Capsule: One capsule daily.

Contraindications

Contraindicated to the patients hypersensitive to any ingredient of the products. Peptic ulcer, hypersensitivity to Diclofenac like other non-steroid anti-inflammatory agents, Diclofenac is also contra-indicated in asthmatic patient in whom attack with asthma, urticaria or acute rhinitis are precipitated by acetylsalicylic acid or by other drugs with prostaglandin synthetase inhibitor.

Side Effects

Moov SR is generally well tolerated. Adverse effects are mild, rare and transient. At the starting of the treatment, however, patients may be sometimes complaining of epigastric pain, eructation, nausea and diarrhea or dizziness or headache. These effects are usually mild in nature. Peripheral edema and skin reactions, such as rash and eczema have also been encountered.

Warning and Precautions

In rare instances where peptic ulceration or gastro-intestinal bleeding occurs in patients under treatment with Diclofenac. In patients with advanced age should be kept under close observation.

Use in pregnancy

During pregnancy Diclofenac should be employed only for compelling reasons. The lowest effective dose should be used. These types of drugs are not recommended during the last trimester of pregnancy.

Use in lactation

Very insignificant quantity of Diclofenac may be detected in breast milk but no undesirable effects on the infant to be expected.

How Supplied

Moov SR Capsule: Each box contains 5X10 capsules in blister pack.

Manufactured by:

 **Astra**
Biopharmaceuticals Ltd.
Gouripur, Ashulia, Dhaka, Bangladesh