

Moov gel

Diclofenac Diethylamine

Composition :

Each gram contains : Diclofenac Diethylamine equivalent to Diclofenac Sodium BP 10 mg.

Descriptive :

Moov gel is an anti-inflammatory and analgesic preparation designed for external use. The white creamy, non-greasy preparation can easily be applied into the skin and its aqueous alcoholic base exerts a soothing and cooling effect on the skin. When applied locally the active substance penetrates the skin, accumulates in the underlying tissue and combats both acute and chronic inflammatory reaction.

The anti-inflammatory and analgesic properties of **Moov gel** elicit a clinical response characterized by a marked decrease in inflammatory swelling of traumatic or rheumatic origin. **Moov gel** affords effective relief from tenderness and pain on movement.

Indications :

For the local treatment of arthritis, periarthropathy, osteoarthritis of peripheral joints and of the vertebral column, localized form of soft-tissue rheumatism syndrome (e.g., tendovaginitis, bursitis, tendonitis), traumatic inflammation of the tendons, ligaments, muscles and joints (e.g., sprains and strain)

Dosage and Applications :

Depending on the size of the area to be treated, 2 - 4 gm of Moov gel should be applied to the skin 3 - 4 times daily to the affected area and applied in lightly. **Moov gel** may also be given in addition to further treatment with other dosage forms of Diclofenac or as prescribed by the physician.

Side Effect :

Moov gel is usually well tolerated. Local irritation, reddening of the skin, skin rash may occasionally occur. When **Moov gel** is applied to a relatively large area of the skin over a prolonged period, the possibility of systemic side-effects cannot be completely excluded.

Contraindication :

Moov gel should not be used to patients hypersensitive to Diclofenac or to the gel base, to acetyl salicylic acid or to other non-steroidal anti-inflammatory agents. **Moov gel** should not be used under occlusive, airtight dressings.

Drug Interaction :

No drug interactions during treatment with diclofenac gel have been reported.

Precautions :

Moov gel should not be allowed to come in contact with the eyes or mucous membranes.

After application, the hands should be washed unless they are the sites being treated. Not to be taken by mouth.

Storage Condition :

Protect from heat and light.

Presentation :

Moov gel 25 gm : Tube containing 25 gm gel.

Manufactured by:

